

High Fluence Iontophoretic Corneal Collagen Cross-linking: In Vivo OCT Imaging of Riboflavin Penetration

To the Editor:

We read with interest the excellent article by Malhotra et al.¹ regarding in vivo estimation of riboflavin penetration using anterior segment optical coherence tomography (OCT). The article evaluates the effect of complete versus grid-like epithelial removal on riboflavin penetration during collagen cross-linking (CXL) in vivo using hand-held OCT. Twenty eyes of 20 patients were imaged intraoperatively at 30 and 60 minutes after starting the procedure. Results showed a homogeneous hyperreflective band extending to a mean depth of 54.2 μm after 30 minutes. In the grid-like removal group, the band was reported uneven in the epi-on areas.

We agree with the authors on the use of OCT for in vivo evaluation of riboflavin's penetration inside the stroma. Moreover, we agree that grid removal of the epithelium is not able to soak the corneal stroma evenly like the epi-off procedure.² In our study, we report the in vivo riboflavin penetration during CXL performed with iontophoresis versus conventional epithelium-off protocol using high-resolution OCT.

Six eyes (6 patients) undergoing CXL were measured preoperatively, intraoperatively, and postoperatively

using high-resolution OCT. The epithelium was removed completely in the central 9-mm zone in 3 eyes (epi-off group), whereas riboflavin penetration through intact epithelium was promoted by an iontophoresis device in the remaining 3 eyes (iontophoresis group). The iontophoresis device for corneal application (8 mm in diameter) is placed on the cornea using an annular suction ring (low suction created by a syringe connected on the suction annulus). The device is filled with approximately 0.5 mL solution from the open proximal side, until the electrode (stainless steel mesh) is covered (**Figure 1A**). The device is connected to a constant current generator (I-ON XL, Sooft, Italy) set at 1 mA (the total dose of 5 mA \times min is monitored by the generator). During the irradiation phase (10mW/cm² for 9 minutes), a good fluorescence is clearly detectable (**Figure 1B**) even after washing with balanced salt solution.

The depth of the hyperreflective band (representing penetration of riboflavin) in the anterior corneal stroma was measured. In the conventional epi-off group, after 30 minutes of passive impregnation, a homogeneous hyperreflective band without fading effect was measured at mean depth of 80 μm (**Figure 1C**). In the iontophoresis group, we observed a less homogeneous but deeper hyperreflective band with a fading effect extending through the anterior 200 μm of the cornea (**Figure 1D**). This band was not visible until the end of irradiation time.

Intraoperative OCT imaging could be a useful technique to evaluate in vivo penetration of riboflavin in-




Figure 1. Iontophoresis impregnation phase (A) throughout an intact epithelium and (B) the irradiation phase in which it is possible to see an intense fluorescence. Intraoperative high-resolution optical coherence tomography images show (C) a homogeneous hyperreflective band at a mean depth of 80 μm in the epi-off group and (D) a less uniform band with a fading effect extending through the anterior 200 μm of the cornea.

side the cornea. Although we still cannot demonstrate the correlation between the band's intensity and riboflavin stromal concentration, we speculate that a higher OCT reflectivity should be positively correlated with it.

These results confirm early pre-clinical evidence that iontophoresis allows penetration of riboflavin through an intact epithelium. According to our hypothesis, the *in vivo* OCT results of the iontophoresis group showed a lower concentration of riboflavin in the stroma compared to the epi-off technique. More studies are needed to evaluate whether this difference could correlate with clinical CXL results.

Paolo Vinciguerra, MD
Miguel Rechichi, MD
Pietro Rosetta, MD
Mario R. Romano, MD, PhD
Leonardo Mastropasqua, MD
Vincenzo Scordia, MD
Claudio Azzolini, MD
Riccardo Vinciguerra, MD
 Istituto Clinico Humanitas
 Milan, Italy

REFERENCES

1. Malhotra C, Shetty R, Kumar RS, et al. *In vivo* imaging of riboflavin penetration during collagen cross-linking with handheld spectral domain optical coherence tomography. *J Refract Surg.* 2012;28:776-780.
2. Samaras K, O'Brart DP, Douth J, et al. Effect of epithelial retention and removal on riboflavin absorption in porcine corneas. *J Refract Surg.* 2009;25:771-775.

Dr. Vinciguerra is a consultant for SOOFT Italy. The remaining authors have no financial or proprietary interest in the materials presented herein.

Reply

We would like to thank Vinciguerra et al. for showing interest in our article.¹ In their study, the authors used iontophoresis to aid riboflavin penetra-

tion during collagen cross-linking and then imaged the effect using a spectral-domain optical coherence tomography. Hao et al. have demonstrated the feasibility of electrically assisted delivery (iontophoresis) of macromolecules into the cornea.² To the best of our knowledge, there have been no published studies on using iontophoresis for riboflavin penetration and we agree with the authors that it provides pre-clinical evidence. The authors might be better served if they could share the device used for the imaging and whether this was done *in vivo* or the subject moved to a device right after the procedure for documenting the changes. As suggested by the authors, larger studies with longer follow-up are required to ascertain both the efficacy and safety of this alternative technique because, based on the authors' results, iontophoresis does not appear to produce adequate or equivocal penetration of riboflavin as the currently used approach.

Harsha Nagaraj, MS
Rohit Shetty, DNB, FRCS (Glasgow)
Rajesh S. Kumar, MS
Himabindu Veluri, MS,
Chintan Malhotra, MS
K. Bhujang Shetty, MS
 Narayana Nethralaya Eye Hospital
 Bangalore, India

REFERENCES

1. Malhotra C, Shetty R, Kumar RS, et al. *In vivo* imaging of riboflavin penetration during collagen cross-linking with handheld spectral domain optical coherence tomography. *J Refract Surg.* 2012;28:776-780.
2. Hao J, Li SK, Liu CY, et al. Electrically assisted delivery of macromolecules into the corneal epithelium. *Exp Eye Res.* 2009;89:934-941.

The authors have no financial or proprietary interest in the materials presented herein.

doi:10.3928/1081597X-20130509-01